### **Condensing and Extracting Against Online Adversaries**



Eshan Chattopadhyay Cornell University



Mohit Gurumukhani Cornell University



Noam Ringach Cornell University



Rocco Servedio Columbia University

**Part 0: Introduction** 

 Useful for randomized algorithms, cryptography, distributed computing protocols, machine learning, etc.

- Useful for randomized algorithms, cryptography, distributed computing protocols, machine learning, etc.
- Most applications need high quality randomness.

- Useful for randomized algorithms, cryptography, distributed computing protocols, machine learning, etc.
- Most applications need high quality randomness.
- In practice, randomness is derived from nature and is of low quality.

### **Extractor**

Weak source (60 / 100)



Uniform source (50 / 50)

### **Extractor**

### Definition (Extractor)

Ext :  $\{0,1\}^n \to \{0,1\}^m$  is  $\varepsilon$ -extractor for class  $\mathcal X$  if for all  $\mathbf X \in \mathcal X$ ,

$$|\mathsf{Ext}(\mathbf{X}) - \mathsf{Uniform}_m| \leq \varepsilon,$$

| · | denotes statistical distance / total variation distance:

$$|\mathbf{A} - \mathbf{B}| = \max_{S \subset \Omega} |\Pr(\mathbf{A} \in S) - \Pr(\mathbf{B} \in S)| = \frac{1}{2} \|\mathbf{A} - \mathbf{B}\|_{1}$$

Single extractor for every

distribution?

Single extractor for every distribution?


## NO!

Single extractor for every distribution?

# NO!

→ Distributions must have entropy

### **Min-entropy**



### Min-entropy

### Definition (Min-entropy)

Min-entropy of source X:

$$H_{\infty}(\mathbf{X}) = -\log\left(\max_{\mathbf{x} \in \mathsf{support}(\mathbf{X})} \Pr(\mathbf{X} = \mathbf{x})\right)$$

### Min-entropy

### Definition (Min-entropy)

Min-entropy of source X:


$$H_{\infty}(\mathbf{X}) = -\log\left(\max_{\mathbf{x} \in \mathsf{support}(\mathbf{X})} \Pr(\mathbf{X} = \mathbf{x})\right)$$

### Definition (Smooth Min-entropy)

Smooth min-entropy of source X with parameter  $\varepsilon$ :

$$H_{\infty}^{\varepsilon}(\mathbf{X}) = \max_{\mathbf{Y}:|\mathbf{X}-\mathbf{Y}|<\varepsilon} H_{\infty}(\mathbf{Y})$$

# NO!



# NO!

Solution: Distributions are structured.

Seedless extractors: a brief history

### Seedless extractors: a brief history

- Two independent sources [Chor-Goldreich'88, ..., Li'23].
- Sources generated by circuits / low complexity classes (applications to circuit lower bounds) [Trevisan Vadhan'00, ..., Viola'14, ...].
- Sources sampled by polynomials over large fields [ Dvir Gabizon Wigderson'09, ... ].
- Sources sampled by polynomials over F₂ [ Chattopadhyay Goodman – Gurumukhani (CGG)'24 ].

Sometimes extractors don't exist

### Sometimes extractors don't exist

• Extractors guarantee closeness to uniform distribution.

### Sometimes extractors don't exist

- Extractors guarantee closeness to uniform distribution.
- Relax this: guarantee closeness to high entropy distribution.

Weak source (60 / 100)



Strong source (48 / 50)

### Condenser

Cond :  $\{0,1\}^n \to \{0,1\}^m$  is a  $(k_{in},k_{out},\varepsilon)$ -condenser for class of distributions  $\mathcal{X}$  with entropy at least  $k_{in}$  if for all  $\mathbf{X} \in \mathcal{X}$ ,

$$H_{\infty}^{\varepsilon}(\mathsf{Cond}(\mathbf{X})) \geq k_{out}$$

### Condenser

Cond :  $\{0,1\}^n \to \{0,1\}^m$  is a  $(k_{in},k_{out},\varepsilon)$ -condenser for class of distributions  $\mathcal{X}$  with entropy at least  $k_{in}$  if for all  $\mathbf{X} \in \mathcal{X}$ ,

$$H_{\infty}^{\varepsilon}(\mathsf{Cond}(\mathbf{X})) \geq k_{out}$$

• Care about increasing entropy rate:

$$\frac{k_{in}}{n}$$
 vs  $\frac{k_{out}}{m}$ 

### Condenser

Cond :  $\{0,1\}^n \to \{0,1\}^m$  is a  $(k_{in}, k_{out}, \varepsilon)$ -condenser for class of distributions  $\mathcal{X}$  with entropy at least  $k_{in}$  if for all  $\mathbf{X} \in \mathcal{X}$ ,

$$H_{\infty}^{\varepsilon}(\mathsf{Cond}(\mathbf{X})) \geq k_{out}$$

Care about increasing entropy rate:

$$\frac{k_{in}}{n} \ll \frac{k_{out}}{m}$$

### Condenser

Cond :  $\{0,1\}^n \to \{0,1\}^m$  is a  $(k_{in},k_{out},\varepsilon)$ -condenser for class of distributions  $\mathcal{X}$  with entropy at least  $k_{in}$  if for all  $\mathbf{X} \in \mathcal{X}$ ,

$$H_{\infty}^{\varepsilon}(\mathsf{Cond}(\mathbf{X})) \geq k_{out}$$

Care about increasing entropy rate:

$$\frac{k_{in}}{n} \ll \frac{k_{out}}{m}$$

• Care about minimizing entropy gap:

$$\Delta_{out} = m - k_{out}$$

Simulating using only weak random source:

Simulating using only weak random source:

• Randomized algorithms with  $poly(\Delta_{out})$  overhead [ Zuckerman'95 ].

Simulating using only weak random source:

- Randomized algorithms with  $\operatorname{poly}(\Delta_{out})$  overhead [ Zuckerman'95 ].
- "One-shot simulations" for randomized protocols, cryptography, interactive proofs etc.

Simulating using only weak random source:

- Randomized algorithms with  $poly(\Delta_{out})$  overhead [ Zuckerman'95 ].
- "One-shot simulations" for randomized protocols, cryptography, interactive proofs etc.

Condensers can exist where extractors (provably) can't

# Single condenser for every high min-entropy distribution?

# Single condenser for every high min-entropy distribution?

# NO!

Single condenser for every high min-entropy distribution?

## NO!

→ Same solution: Distributions should be structured.

Seedless condensers: prior work

# Seedless condensers: prior work

- Condensers for Chor-Goldreich (CG) sources and adversarial
  Chor-Coldreich (CG) sources [ Doron Moshkovitz Oh Zuckerman'23 ].
- Improved Condensers for Chor-Goldreich Sources [Goodman Li Zuckerman'24]

**Part 1: Models and Results** 

### Oblivious Bit Fixing Sources (OBFs)

•  $\ell$  bit input.



### Oblivious Bit Fixing Sources (OBFs)

- ullet bit input.
- g good bits: uniform,  $\ell-g$  bad bits: constants.



### Oblivious Bit Fixing Sources (OBFs)

- $\ell$  bit input.
- g good bits: uniform,  $\ell-g$  bad bits: constants.




PARITY extracts from  $(1, \ell)$ -OBFs.

# **NOBFs**

### **NOBFs**

### Non-Oblivious Bit Fixing Sources (NOBFs)

• g good bits: uniform,  $\ell-g$  bad bits: arbitrary functions of good bits.



Kahn-Kalai-Linial'88, Ben-Or-Linial'89, Ajtai-Linial'93

• Can't extract from  $\left(\ell - \frac{\ell}{\log(\ell)}, \ell\right)$ -NOBFs.

#### Kahn - Kalai - Linial'88, Ben-Or - Linial'89, Ajtai - Linial'93

- Can't extract from  $\left(\ell \frac{\ell}{\log(\ell)}, \ell\right)$ -NOBFs.
- Can extract from  $\left(\ell \frac{\ell}{\log^2(\ell)}, \ell\right)$ -NOBFs.

#### Kahn-Kalai-Linial'88, Ben-Or-Linial'89, Ajtai-Linial'93

- Can't extract from  $\left(\ell \frac{\ell}{\log(\ell)}, \ell\right)$ -NOBFs.
- Can extract from  $\left(\ell \frac{\ell}{\log^2(\ell)}, \ell\right)$ -NOBFs.

#### **Ouestion**

Can you condense from  $(g,\ell)$ -NOBFs when  $g<\ell-rac{\ell}{\log\ell}$ ?

Kahn - Kalai - Linial'88, Ben-Or - Linial'89, Ajtai - Linial'93

- Can't extract from  $\left(\ell \frac{\ell}{\log(\ell)}, \ell\right)$ -NOBFs.
- Can extract from  $\left(\ell \frac{\ell}{\log^2(\ell)}, \ell\right)$ -NOBFs.

### Theorem (Chattopadhyay - Gurumukhani - R (CGR)'24)

For constant  $\alpha$ , **can't** condense  $((1-\alpha)\cdot \ell,\ell)$ -NOBFs beyond **rate 1 — \alpha**.

# **NOSFs**

### **NOSFs**


### Non-Oblivious Symbol Fixing Sources (NOSFs)

•  $\ell$  blocks of length n each.

### **NOSFs**

### Non-Oblivious Symbol Fixing Sources (NOSFs)

- $\ell$  blocks of length n each.
- g good blocks: uniform,  $\ell-g$  bad blocks: arbitrary functions of good blocks.



Aggarwal – Obremski – Ribeiro – Siniscalchi – Visconti (AORSV)'20

**Can't** extract from  $(0.99\ell, \ell)$ -NOSFs.

Aggarwal – Obremski – Ribeiro – Siniscalchi – Visconti (AORSV)'20

**Can't** extract from  $(0.99\ell,\ell)$ -NOSFs.

#### Question

Can you condense from  $(g, \ell)$ -NOSFs?

Aggarwal – Obremski – Ribeiro – Siniscalchi – Visconti (AORSV)'20

**Can't** extract from  $(0.99\ell,\ell)$ -NOSFs.

#### Theorem (CGR'24)

**Can't** condense  $(g, \ell)$ -NOSFs beyond **rate**  $g/\ell$ .

# **oNOSFs**

### **oNOSFs**


### Online Non-Oblivious Symbol Fixing Sources (oNOSFs)

• g good blocks: uniform,  $\ell-g$  bad blocks: arbitrary functions of good blocks **that appear before it**.


### **oNOSFs**

### Online Non-Oblivious Symbol Fixing Sources (oNOSFs)


• g good blocks: uniform,  $\ell-g$  bad blocks: arbitrary functions of good blocks **that appear before it**.
















[AORSV'20]

**Can't** extract from  $(0.99\ell, \ell)$ -oNOSFs.

### [AORSV'20]

**Can't** extract from  $(0.99\ell,\ell)$ -oNOSFs.

#### Question

Can you condense from  $(g, \ell)$ -oNOSFs?

### [AORSV'20]

**Can't** extract from  $(0.99\ell, \ell)$ -oNOSFs.

### Theorem (CGR'24, CGRS'25)


Can't condense  $(g, \ell)$ -oNOSFs beyond rate  $\frac{1}{\lfloor \ell/g \rfloor}$ .

### [AORSV'20]

**Can't** extract from  $(0.99\ell, \ell)$ -oNOSFs.

### Theorem (CGR'24, CGRS'25)

Can't condense  $(g,\ell)$ -oNOSFs beyond rate  $\frac{1}{\lfloor \ell/g \rfloor}$ . Can condense  $(g,\ell)$ -oNOSFs to rate  $\frac{1}{\lfloor \ell/g \rfloor}$ .



# **Extracting / Condensing from oNOSFs**

Corollary (Sharp threshold at  $\boldsymbol{g}=\ell/2$ )

**Can't** condense  $(0.5\ell,\ell)$ -oNOSFs beyond **rate** 1/2 - Impossibility.

**Can** condense  $(0.51\ell,\ell)$ -oNOSFs to **rate** 0.99 - Possibility.

**Part 2: Possibility** 

#### Theorem (Condensing uniform oNOSF sources)

For  $g \ge 0.51\ell$ , large constant block length n, and  $\ell$  increasing, we can condense any oNOSF source to entropy rate 0.99.

#### Theorem (Condensing uniform oNOSF sources)

```
For g \geq 0.51\ell, \ell = \Omega(\log(1/\varepsilon)), and n = 10^4, exists \mathrm{Cond}: \{0,1\}^{\ell n} \to \{0,1\}^m s.t. for any (g,\ell)-oNOSF \mathbf{X}, \mathbf{H}_{\infty}^{\varepsilon}(\mathrm{Cond}(\mathbf{X})) \geq 0.99m where m = \Omega(\ell + \log(1/\varepsilon)).
```

#### Theorem (Condensing uniform oNOSF sources)

For  $g \geq 0.51\ell$ ,  $\ell = \Omega(\log(1/\varepsilon))$ , and  $n = 10^4$ , exists  $\mathrm{Cond}: \{0,1\}^{\ell n} \to \{0,1\}^m$  s.t. for any  $(g,\ell)$ -oNOSF  $\mathbf{X}$ ,  $\mathbf{H}^{\varepsilon}_{\infty}(\mathrm{Cond}(\mathbf{X})) \geq 0.99m$  where  $m = \Omega(\ell + \log(1/\varepsilon))$ .

#### Theorem (Condensing low-entropy oNOSF sources)

For  $g \ge 0.51\ell$ , we can similarly condense oNOSF sources with logarithmic min-entropy.

#### Theorem (Condensing uniform oNOSF sources)

For  $g \geq 0.51\ell$ ,  $\ell = \Omega(\log(1/\varepsilon))$ , and  $n = 10^4$ , exists  $\mathrm{Cond}: \{0,1\}^{\ell n} \to \{0,1\}^m$  s.t. for any  $(g,\ell)$ -oNOSF  $\mathbf{X}$ ,  $\mathbf{H}^{\varepsilon}_{\infty}(\mathrm{Cond}(\mathbf{X})) \geq 0.99m$  where  $m = \Omega(\ell + \log(1/\varepsilon))$ .

#### Theorem (Condensing low-entropy oNOSF sources)

For  $g \geq 0.51\ell$ ,  $n = \operatorname{polylog}(\ell/\varepsilon)$  exists  $\operatorname{Cond}: (\{0,1\}^n)^\ell \to \{0,1\}^m$  s.t. for any low-entropy  $(g,\ell)$ -oNOSF  $\mathbf X$  with  $k = \Omega(\log(\ell/\varepsilon))$ ,  $H^\varepsilon_\infty(\operatorname{Cond}(\mathbf X)) \geq m - \operatorname{O}(m/\log m) - \operatorname{O}(\log(1/\varepsilon))$  where  $m = \Omega(k)$ .

#### Theorem (Condensing uniform oNOSF sources)

For  $g \geq 0.51\ell$ ,  $\ell = \Omega(\log(1/\varepsilon))$ , and  $n = 10^4$ , exists  $\mathrm{Cond}: \{0,1\}^{\ell n} \to \{0,1\}^m$  s.t. for any  $(g,\ell)$ -oNOSF  $\mathbf{X}$ ,  $\mathbf{H}^\varepsilon_\infty(\mathbf{Cond}(\mathbf{X})) \geq 0.99m$  where  $m = \Omega(\ell + \log(1/\varepsilon))$ .

#### Theorem (Condensing low-entropy oNOSF sources)

For  $g \geq 0.51\ell$ ,  $n = \text{polylog}(\ell/\varepsilon)$  exists Cond :  $(\{0,1\}^n)^\ell \to \{0,1\}^m$  s.t. for any low-entropy  $(g,\ell)$ -oNOSF X with  $k = \Omega(\log(\ell/\varepsilon))$ ,  $H_{\infty}^{\epsilon}(\text{Cond}(X)) \geq m - O(m/\log m) - O(\log(1/\varepsilon))$  where  $m = \Omega(k)$ .

#### Theorem (Extend AORSV'20 result)

Transform low-entropy  $(g,\ell)$ -oNOSFs  $\rightarrow$  uniform  $(0.99g,\ell)$ -oNOSFs.

#### Theorem (Condensing uniform oNOSF sources)

```
For g \geq 0.51\ell, \ell = \Omega(\log(1/\varepsilon)), and n = 10^4, exists Cond : \{0,1\}^{\ell n} \to \{0,1\}^m s.t. for any (g,\ell)-oNOSF \mathbf{X}, \mathbf{H}_{\infty}^{\varepsilon}(\mathbf{Cond}(\mathbf{X})) \geq \mathbf{0.99m} where m = \Omega(\ell + \log(1/\varepsilon)).
```

# Does a random function work?

#### Theorem (Condensing uniform oNOSF sources)


```
For g \geq 0.51\ell, \ell = \Omega(\log(1/\varepsilon)), and n = 10^4, exists \mathrm{Cond}: \{0,1\}^{\ell n} \to \{0,1\}^m s.t. for any (g,\ell)-oNOSF \mathbf{X}, \mathbf{H}^{\varepsilon}_{\infty}(\mathrm{Cond}(\mathbf{X})) \geq 0.99m where m = \Omega(\ell + \log(1/\varepsilon)).
```

# Does a random function work?

# No!


#### Theorem (Condensing uniform oNOSF sources)

For  $g \geq 0.51\ell$ ,  $\ell = \Omega(\log(1/\varepsilon))$ , and  $n = 10^4$ , exists Cond :  $\{0,1\}^{\ell n} \to \{0,1\}^m$  s.t. for any  $(g,\ell)$ -oNOSF  $\mathbf{X}$ ,  $\mathbf{H}^{\varepsilon}_{\infty}(\mathbf{Cond}(\mathbf{X})) \geq \mathbf{0.99m}$  where  $m = \Omega(\ell + \log(1/\varepsilon))$ .



#### Theorem (Condensing uniform oNOSF sources)

For  $g \geq 0.51\ell$ ,  $\ell = \Omega(\log(1/\varepsilon))$ , and  $n = 10^4$ , exists Cond :  $\{0,1\}^{\ell n} \to \{0,1\}^m$  s.t. for any  $(g,\ell)$ -oNOSF  $\mathbf{X}$ ,  $\mathbf{H}_{\infty}^{\varepsilon}(\mathbf{Cond}(\mathbf{X})) \geq \mathbf{0.99m}$  where  $m = \Omega(\ell + \log(1/\varepsilon))$ .



#### Problem

A random function doesn't condense because the adversary has too much power in latter blocks.





#### Solution


Take only first bit of latter half of blocks to weaken the adversary.






#### Solution


Take only first bit of latter half of blocks to weaken the adversary.






#### Solution

Take only first bit of latter half of blocks to weaken the adversary.







# Now a random function works!

#### Solution

Take only first bit of latter half of blocks to weaken the adversary.





#### Formal definition

A function sCond :  $\{0,1\}^n \times \{0,1\}^d \to \{0,1\}^m$  is a  $(k,\varepsilon)$ -seeded condenser for a class of sources  $\mathcal{X}$  if for all  $\mathbf{X} \in \mathcal{X}$ ,

$$H_{\infty}^{\varepsilon}(\mathsf{sCond}(\mathbf{X},\mathbf{U}_d)) \geq k$$

#### Formal definition

A function sCond :  $\{0,1\}^n \times \{0,1\}^d \to \{0,1\}^m$  is a  $(k,\varepsilon)$ -seeded condenser for a class of sources  $\mathcal{X}$  if for all  $\mathbf{X} \in \mathcal{X}$ ,

$$H_{\infty}^{\varepsilon}(\mathsf{sCond}(\mathbf{X},\mathbf{U}_d)) \geq k$$

#### Theorem (Good seeded condensers exist)

Seeded condensers with logarithmic seed length and linear output length exist.

#### Formal definition


A function sCond :  $\{0,1\}^n \times \{0,1\}^d \to \{0,1\}^m$  is a  $(k,\varepsilon)$ -seeded condenser for a class of sources  $\mathcal{X}$  if for all  $\mathbf{X} \in \mathcal{X}$ ,

$$H_{\infty}^{\varepsilon}(\mathsf{sCond}(\mathbf{X},\mathbf{U}_d)) \geq k$$

#### Theorem (Good seeded condensers exist)


For all  $d, \varepsilon$  s.t.  $d \ge \log(\ell n/\varepsilon) + O(1)$  and  $m = 0.01\ell n + d + \log(1/\varepsilon) + O(1)$ , exists sCond :  $\{0,1\}^{\ell n/2} \times \{0,1\}^d \to \{0,1\}^m$  s.t. for all  $\mathbf{X} \sim \{0,1\}^{\ell n/2}$  with  $H_{\infty}(\mathbf{X}) \ge 0.01\ell n$ , we have

$$H_{\infty}^{\varepsilon}(\mathsf{sCond}(\mathbf{X}, \mathbf{U}_d)) \geq 0.01 \ell n + d$$




#### Proof.


•  $g \geq 0.51\ell \implies$  at least  $0.01\ell$  blocks in  $\mathbf{Z}_1 \sim \{0,1\}^{\ell n/2}$  are good.




- $g \ge 0.51\ell \implies$  at least  $0.01\ell$  blocks in  $\mathbf{Z}_1 \sim \{0,1\}^{\ell n/2}$  are good.
- Pretend  $\mathbf{Z}_2 \sim \{0,1\}^{\ell/2}$  is uniform.




- $g \ge 0.51\ell \implies$  at least  $0.01\ell$  blocks in  $\mathbf{Z}_1 \sim \{0,1\}^{\ell n/2}$  are good.
- Pretend  $\mathbf{Z}_2 \sim \{0,1\}^{\ell/2}$  is uniform.
- sCond requirements satisfied!



- $g \ge 0.51\ell \implies$  at least  $0.01\ell$  blocks in  $\mathbf{Z}_1 \sim \{0,1\}^{\ell n/2}$  are good.
- Pretend  $\mathbf{Z}_2 \sim \{0,1\}^{\ell/2}$  is uniform.
- sCond requirements satisfied!
- ullet  $H_{\infty}^{arepsilon_{ ext{sCond}}}( ext{sCond}(\mathbf{Z}_1,\mathbf{U}_{\ell/2})) \geq k_{ ext{sCond}}.$



- $g \ge 0.51\ell \implies$  at least  $0.01\ell$  blocks in  $\mathbf{Z}_1 \sim \{0,1\}^{\ell n/2}$  are good.
- Pretend  $\mathbf{Z}_2 \sim \{0,1\}^{\ell/2}$  is uniform.
- sCond requirements satisfied!
- $H_{\infty}^{\varepsilon_{\mathsf{sCond}}}(\mathsf{sCond}(\mathbf{Z}_1,\mathbf{U}_{\ell/2})) \geq k_{\mathsf{sCond}}.$
- BUT  $\mathbf{Z}_2$  might have  $0.49\ell$  bad bits!



Adversary can't make things too bad

# Adversary can't make things too bad


#### Lemma

Let  $\mathbf{X} \sim \{0,1\}^n$  and sCond be s.t.  $H_{\infty}^{\epsilon_{\mathsf{sCond}}}(\mathsf{sCond}(\mathbf{X},\mathbf{U}_d)) \geq k_{\mathsf{sCond}}$ . Let  $\mathbf{U}_d'$  be  $\mathbf{U}_d$  except an adversary controls some b bits. Then,


$$H_{\infty}^{\varepsilon'}(\mathsf{sCond}(\mathbf{X},\mathbf{U}_d')) \geq k_{\mathsf{sCond}} - b$$

where  $\varepsilon' = \varepsilon_{sCond} \cdot 2^b$ .


- $g \ge 0.51\ell \implies$  at least  $0.01\ell$  blocks in  $\mathbf{Z}_1 \sim \{0,1\}^{\ell n/2}$  are good.
- Pretend  $\mathbf{Z}_2 \sim \{0,1\}^{\ell/2}$  is uniform.
- sCond requirements satisfied!
- $H_{\infty}^{\varepsilon_{\mathsf{sCond}}}(\mathsf{sCond}(\mathbf{Z}_1,\mathbf{U}_{\ell/2})) \geq k_{\mathsf{sCond}}$ .
- BUT  $\mathbf{Z}_2$  might have  $0.49\ell$  bad bits!



- $g \ge 0.51\ell \implies$  at least  $0.01\ell$  blocks in  $\mathbf{Z}_1 \sim \{0,1\}^{\ell n/2}$  are good.
- Pretend  $\mathbf{Z}_2 \sim \{0,1\}^{\ell/2}$  is uniform.
- sCond requirements satisfied!
- ullet  $H_{\infty}^{arepsilon_{ ext{sCond}}}( ext{sCond}(\mathbf{Z}_1,\mathbf{U}_{\ell/2})) \geq k_{ ext{sCond}}.$
- BUT  $\mathbf{Z}_2$  might have  $0.49\ell$  bad bits!
- Save sCond using lemma that adversary can't be too bad.



- $g \ge 0.51\ell \implies$  at least  $0.01\ell$  blocks in  $\mathbf{Z}_1 \sim \{0,1\}^{\ell n/2}$  are good.
- Pretend  $\mathbf{Z}_2 \sim \{0,1\}^{\ell/2}$  is uniform.
- sCond requirements satisfied!
- $H_{\infty}^{\varepsilon_{\mathsf{sCond}}}(\mathsf{sCond}(\mathbf{Z}_1,\mathbf{U}_{\ell/2})) \geq k_{\mathsf{sCond}}.$
- BUT  $\mathbf{Z}_2$  might have  $0.49\ell$  bad bits!
- Save sCond using lemma that adversary can't be too bad.
- $H_{\infty}^{\varepsilon'}(\operatorname{sCond}(\mathbf{Z}_1, \mathbf{Z}_2) \ge k_{\operatorname{sCond}} 0.49\ell$ with  $\varepsilon' = \varepsilon_{\operatorname{sCond}} \cdot 2^{0.49\ell}$ .



#### Proof.

- $g \ge 0.51\ell \implies$  at least  $0.01\ell$  blocks in  $\mathbf{Z}_1 \sim \{0,1\}^{\ell n/2}$  are good.
- Pretend  $\mathbf{Z}_2 \sim \{0,1\}^{\ell/2}$  is uniform.
- sCond requirements satisfied!
- $H_{\infty}^{\varepsilon_{\mathsf{sCond}}}(\mathsf{sCond}(\mathbf{Z}_1,\mathbf{U}_{\ell/2})) \geq k_{\mathsf{sCond}}$ .
- BUT  $\mathbf{Z}_2$  might have  $0.49\ell$  bad bits!
- Save sCond using lemma that adversary can't be too bad.
- $H_{\infty}^{\varepsilon'}(\operatorname{sCond}(\mathbf{Z}_1, \mathbf{Z}_2) \geq k_{\operatorname{sCond}} 0.49\ell$ with  $\varepsilon' = \varepsilon_{\operatorname{sCond}} \cdot 2^{0.49\ell}$ .

#### Proof.

• To get error  $\varepsilon$ , need  $\varepsilon_{\text{scond}} = \varepsilon \cdot 2^{-0.49\ell}$ .

#### Proof.

- $g \ge 0.51\ell \implies$  at least  $0.01\ell$  blocks in  $\mathbf{Z}_1 \sim \{0,1\}^{\ell n/2}$  are good.
- Pretend  $\mathbf{Z}_2 \sim \{0,1\}^{\ell/2}$  is uniform.
- sCond requirements satisfied!
- ullet  $H_{\infty}^{arepsilon_{ ext{sCond}}}( ext{sCond}(\mathbf{Z}_1,\mathbf{U}_{\ell/2})) \geq k_{ ext{sCond}}.$
- BUT  $\mathbf{Z}_2$  might have  $0.49\ell$  bad bits!
- Save sCond using lemma that adversary can't be too bad.
- $H_{\infty}^{\varepsilon'}(\operatorname{sCond}(\mathbf{Z}_1, \mathbf{Z}_2) \geq k_{\operatorname{sCond}} 0.49\ell$ with  $\varepsilon' = \varepsilon_{\operatorname{sCond}} \cdot 2^{0.49\ell}$ .

- To get error  $\varepsilon$ , need  $\varepsilon_{\mathsf{sCond}} = \varepsilon \cdot 2^{-0.49\ell}$ .
- For sCond to exist, need  $0.5\ell > \log(\ell n/\varepsilon_{\text{sCond}}) + O(1)$

#### Proof.

- $g \ge 0.51\ell \implies$  at least  $0.01\ell$  blocks in  $\mathbf{Z}_1 \sim \{0,1\}^{\ell n/2}$  are good.
- Pretend  $\mathbf{Z}_2 \sim \{0,1\}^{\ell/2}$  is uniform.
- sCond requirements satisfied!
- $H_{\infty}^{\varepsilon_{\mathsf{sCond}}}(\mathsf{sCond}(\mathbf{Z}_1,\mathbf{U}_{\ell/2})) \geq k_{\mathsf{sCond}}.$
- BUT  $\mathbf{Z}_2$  might have  $0.49\ell$  bad bits!
- Save sCond using lemma that adversary can't be too bad.
- $H_{\infty}^{\varepsilon'}(\operatorname{sCond}(\mathbf{Z}_1, \mathbf{Z}_2) \geq k_{\operatorname{sCond}} 0.49\ell$ with  $\varepsilon' = \varepsilon_{\operatorname{sCond}} \cdot 2^{0.49\ell}$ .

- To get error  $\varepsilon$ , need  $\varepsilon_{\mathsf{sCond}} = \varepsilon \cdot 2^{-0.49\ell}$ .
- For sCond to exist, need  $0.5\ell \geq \log(\ell n/\varepsilon_{\rm sCond}) + {\it O}(1)$
- So require  $\varepsilon \geq 2^{-0.01\ell}$ .

#### Proof.

- $g \ge 0.51\ell \implies$  at least  $0.01\ell$  blocks in  $\mathbf{Z}_1 \sim \{0,1\}^{\ell n/2}$  are good.
- Pretend  $\mathbf{Z}_2 \sim \{0,1\}^{\ell/2}$  is uniform.
- sCond requirements satisfied!
- $H^{\varepsilon_{\mathsf{sCond}}}_{\infty}(\mathsf{sCond}(\mathbf{Z}_1,\mathbf{U}_{\ell/2})) \geq k_{\mathsf{sCond}}$ .
- BUT  $\mathbf{Z}_2$  might have  $0.49\ell$  bad bits!
- Save sCond using lemma that adversary can't be too bad.
- $H_{\infty}^{\varepsilon'}(\operatorname{sCond}(\mathbf{Z}_1, \mathbf{Z}_2) \ge k_{\operatorname{sCond}} 0.49\ell$ with  $\varepsilon' = \varepsilon_{\operatorname{sCond}} \cdot 2^{0.49\ell}$ .

- To get error  $\varepsilon$ , need  $\varepsilon_{\mathsf{sCond}} = \varepsilon \cdot 2^{-0.49\ell}$ .
- For sCond to exist, need  $0.5\ell \ge \log(\ell n/\varepsilon_{\text{sCond}}) + O(1)$
- So require  $\varepsilon > 2^{-0.01\ell}$ .
- sCond outputs  $m = 0.01 \ell n + O(1)$  bits with entropy  $m O(\ell)$ .

#### Proof.

- $g \ge 0.51\ell \implies$  at least  $0.01\ell$  blocks in  $\mathbf{Z}_1 \sim \{0,1\}^{\ell n/2}$  are good.
- Pretend  $\mathbf{Z}_2 \sim \{0,1\}^{\ell/2}$  is uniform.
- sCond requirements satisfied!
- $H_{\infty}^{\varepsilon_{\mathsf{sCond}}}(\mathsf{sCond}(\mathbf{Z}_1,\mathbf{U}_{\ell/2})) \geq k_{\mathsf{sCond}}$ .
- BUT  $\mathbf{Z}_2$  might have  $0.49\ell$  bad bits!
- Save sCond using lemma that adversary can't be too bad.
- $H_{\infty}^{\varepsilon'}(\operatorname{sCond}(\mathbf{Z}_1, \mathbf{Z}_2) \ge k_{\operatorname{sCond}} 0.49\ell$ with  $\varepsilon' = \varepsilon_{\operatorname{sCond}} \cdot 2^{0.49\ell}$ .

- To get error  $\varepsilon$ , need  $\varepsilon_{\mathsf{sCond}} = \varepsilon \cdot 2^{-0.49\ell}$ .
- For sCond to exist, need  $0.5\ell \ge \log(\ell n/\varepsilon_{\text{sCond}}) + O(1)$
- So require  $\varepsilon > 2^{-0.01\ell}$ .
- sCond outputs  $m = 0.01 \ell n + O(1)$  bits with entropy  $m O(\ell)$ .
- So for large const n, get entropy rate  $\frac{m-O(\ell)}{m} \ge 0.99$ .

# **Condensing from oNOSFs**

#### Corollary (Sharp threshold at $m{g}=\ell/2$ )

**Can't** condense  $(\ell/2,\ell)$ -oNOSFs beyond **rate** 1/2 - Impossibility.

**Can** condense  $(0.51\ell,\ell)$ -oNOSFs to **rate** 0.99 - Possibility.  $\checkmark$ 

# Part 3: Future Directions

• Explicit condensers for oNOSFs with constant block length?

- Explicit condensers for oNOSFs with constant block length?
- Our condensers (explicit and existential) have super-constant output entropy gap. Is constant possible (for any values of  $\ell$  and n)?

- Explicit condensers for oNOSFs with constant block length?
- Our condensers (explicit and existential) have super-constant output entropy gap. Is constant possible (for any values of  $\ell$  and n)?
- Construct condensers for oNOBFs (online NOBFs).

- Explicit condensers for oNOSFs with constant block length?
- Our condensers (explicit and existential) have super-constant output entropy gap. Is constant possible (for any values of  $\ell$  and n)?
- Construct condensers for oNOBFs (online NOBFs).

